

Request for Prior Authorization for Alpha-1 Proteinase Inhibitors Website Form – <u>www.wv.highmarkhealthoptions.com</u> Submit request via: Fax - 1-833-547-2030

All requests for Alpha-1 Proteinase Inhibitors require a Prior Authorization and will be screened for medical necessity and appropriateness using the criteria listed below.

Alpha-1 Proteinase Inhibitors include Aralast NP, Glassia, Prolastin-C and Zemaira. New products with this classification will require the same documentation.

## Alpha-1 Proteinase Inhibitors Prior Authorization Criteria:

Coverage may be provided with a <u>diagnosis</u> of Emphysema due to congenital deficiency of alpha1proteinase inhibitor (A1-PI) and the following criteria is met:

- Member has a diagnosis of congenital alpha-1-antitrypsin deficiency (AATD) confirmed by **ONE** of the following:
  - A high risk AATD genetic variant [e.g., Pi\*ZZ, Pi\*Z(null), Pi\*(null)(null), or Pi\*SZ protein phenotypes (homozygous)]
  - $\circ~$  Other rare AAT deficiency disease-causing alleles associated with serum AAT level < 11  $\mu mol/L$
- Member has a baseline circulating serum concentration of AATD  $< 11 \mu$ mol/L using rocket immunoelectrophoresis (which corresponds to < 80 mg/dl if measured by radial immunodiffusion or < 57 mg/dl if measured by nephelometry).
- Member has a diagnosis of clinically evident emphysema confirmed by **ONE** of the following:
  - Forced expiratory volume in one second (FEV1) from  $\ge$  30% to  $\le$  65% of predicted, post-bronchodilator
  - $\circ~$  FEV1 from > 65% to < 80% of predicted, post-bronchodilator, and a rapid decline in lung function showing a change in FEV1 > 100 mL/year
- Medication is prescribed by or in consultation with a pulmonologist.
- Prescriber attests that member will continue to be on optimal conventional treatment for emphysema (e.g., bronchodilators, supplemental oxygen, etc.)
- Member is currently a nonsmoker or ex-smoker
- Member must not have a contraindication to therapy such as an Immunoglobulin A (IgA) deficiency with antibodies against IgA.
- Must be age-appropriate according to FDA-approved labeling, nationally recognized compendia, or evidence-based practice guidelines
- The requested dose and frequency is in accordance with FDA-approved labeling, nationally recognized compendia, and/or evidence-based practice guidelines
- Initial Duration of Approval: 6 months
- Reauthorization criteria
  - Reauthorization benefit will be approved if there is documentation of improvement or stabilization of the signs and symptoms of emphysema associated with alpha-1



Updated: 06/2024 Approved: 06/2024

antitrypsin deficiency including slowed progression of emphysema as evidenced by annual spirometry testing or a decrease in frequency, duration or severity of pulmonary exacerbations

## • Reauthorization Duration of Approval: 12 months

Coverage may be provided for any non-FDA labeled indication if it is determined that the use is a medically accepted indication supported by nationally recognized pharmacy compendia or peer-reviewed medical literature for treatment of the diagnosis(es) for which it is prescribed. These requests will be reviewed on a case by case basis to determine medical necessity.

Drugs are authorized in generic form unless the branded product is on the preferred drug list or the prescriber has indicated in writing that the branded product is medically necessary. If only the branded product is on the preferred drug list, the generic form will be considered non-preferred and shall not require the prescriber to indicate in writing that the branded product is medically necessary.



Updated: 06/2024 Approved: 06/2024

ALPHA-1 PROTEINASE INHIBITORS PRIOR AUTHORIZATION FORM				
Please complete and fax all requested information below including any progress notes, laboratory test results, or chart documentation				
as applicable to Highmark Health Options Pharmacy Services. FAX: (833)-547-2030				
			: (844) 325-6251 Mon – Fri 8 am to 7 pm	
PROVIDER INFORMATION				
Requesting Provider:		NPI:		
Provider Specialty: Offi		Office C	ontact:	
ffice Address: Offi		Office Pl	one:	
		Office Fa	х:	
MEMBER INFORMATION				
Member Name:		DOB:		
Member ID:		Member weight:	Height:	
REQUESTED DRUG INFORMATION				
Medication:		Strength:		
Directions:		Quantity:	Refills:	
Is the member currently receiving rec			Medication Initiated:	
Is this medication being used for a chronic or long-term condition for which the medication may be necessary for the life of the				
patient? Yes No				
Billing Information				
	<u> </u>	ally, JCODE:		
Place of Service: Hospital Provider's office Member's home Other				
Num	Place of Serv	ice Information		
Name:		NPI:		
Address:		Phone:		
	MEDICAL HISTORY (Complete for ALL requests)			
Diagnosis: Emphysema due to co				
	ongenital deficiency of A-1	PI		
Does the member have a diagnosis of	ongenital deficiency of A-1 f congenital alpha-1-antitryp	PI	) confirmed by ONE of the following?	
Does the member have a diagnosis of	ongenital deficiency of A-1 f congenital alpha-1-antitryp	PI		
Does the member have a diagnosis of • A high risk AATD genetic v	ongenital deficiency of A-1 f congenital alpha-1-antitryp variant [e.g., Pi*ZZ, Pi*Z(nu	PI D Other: sin deficiency (AATI II), Pi*(null)(null), or	) confirmed by ONE of the following? Pi*SZ protein phenotypes (homozygous)]	
<ul> <li>Does the member have a diagnosis of</li> <li>A high risk AATD genetic v Yes □ No</li> </ul>	ongenital deficiency of A-1 f congenital alpha-1-antitryp variant [e.g., Pi*ZZ, Pi*Z(nu ausing alleles associated with	PI DOther: sin deficiency (AATE III), Pi*(null)(null), or h serum AAT level <	) confirmed by ONE of the following? Pi*SZ protein phenotypes (homozygous)]	
<ul> <li>Does the member have a diagnosis of</li> <li>A high risk AATD genetic v Yes No</li> <li>Other rare AATD disease-ca</li> <li>What is the member's baseline circula</li> <li>Does the member have any contraind</li> </ul>	ongenital deficiency of A-1 f congenital alpha-1-antitryp variant [e.g., Pi*ZZ, Pi*Z(nu ausing alleles associated with ating serum concentration of ications to therapy? Yes	PI Other: sin deficiency (AATI Ill), Pi*(null)(null), or h serum AAT level < AATD? No	) confirmed by ONE of the following? Pi*SZ protein phenotypes (homozygous)]	
<ul> <li>Does the member have a diagnosis of</li> <li>A high risk AATD genetic v Yes No</li> <li>Other rare AATD disease-ca</li> <li>What is the member's baseline circula</li> <li>Does the member have any contrained</li> <li>Does the member have a diagnosis of</li> </ul>	ongenital deficiency of A-1 f congenital alpha-1-antitryp variant [e.g., Pi*ZZ, Pi*Z(nu ausing alleles associated with ating serum concentration of lications to therapy? Yes f emphysema confirmed by a	PI 🗍 Other: sin deficiency (AATI ill), Pi*(null)(null), or h serum AAT level < f AATD? No any of the following?	) confirmed by ONE of the following? Pi*SZ protein phenotypes (homozygous)] 11 μmol/L Yes Νο	
<ul> <li>Does the member have a diagnosis of</li> <li>A high risk AATD genetic v Yes No</li> <li>Other rare AATD disease-ca</li> <li>What is the member's baseline circula</li> <li>Does the member have any contraind</li> </ul>	ongenital deficiency of A-1 f congenital alpha-1-antitryp variant [e.g., Pi*ZZ, Pi*Z(nu ausing alleles associated with ating serum concentration of lications to therapy? Yes f emphysema confirmed by a	PI 🗍 Other: sin deficiency (AATI ill), Pi*(null)(null), or h serum AAT level < f AATD? No any of the following?	) confirmed by ONE of the following? Pi*SZ protein phenotypes (homozygous)] 11 μmol/L Yes Νο	
<ul> <li>Does the member have a diagnosis of</li> <li>A high risk AATD genetic v Yes No</li> <li>Other rare AATD disease-ca</li> <li>What is the member's baseline circula</li> <li>Does the member have any contraind</li> <li>Does the member have a diagnosis of</li> <li>Forced expiratory volume in</li> </ul>	ongenital deficiency of A-1 f f congenital alpha-1-antitryp variant [e.g., Pi*ZZ, Pi*Z(nu ausing alleles associated with ating serum concentration of fications to therapy? $\Box$ Yes f emphysema confirmed by a n one second (FEV1) from $\geq$	PI $\Box$ Other: sin deficiency (AATI ill), Pi*(null)(null), or h serum AAT level < f AATD? $\Box$ No any of the following? 30% to $\leq 65\%$ of pre-	) confirmed by ONE of the following? Pi*SZ protein phenotypes (homozygous)] 11 μmol/L Yes Νο	
<ul> <li>Does the member have a diagnosis of</li> <li>A high risk AATD genetic v Yes □ No</li> <li>Other rare AATD disease-ca</li> <li>What is the member's baseline circula</li> <li>Does the member have any contraind</li> <li>Does the member have a diagnosis of</li> <li>Forced expiratory volume in</li> <li>FEV1 from &gt; 65% to &lt; 80% &gt; 100 mL/year</li> </ul>	ongenital deficiency of A-1 f f congenital alpha-1-antitryp variant [e.g., Pi*ZZ, Pi*Z(nu ausing alleles associated with ating serum concentration of fications to therapy? $\Box$ Yes f emphysema confirmed by a n one second (FEV1) from $\geq$ of predicted, post-bronchodi	PI $\Box$ Other: sin deficiency (AATE ill), Pi*(null)(null), or h serum AAT level < f AATD? $\Box$ No any of the following? 30% to $\leq 65\%$ of pre- ilator, and a rapid decl	) confirmed by ONE of the following? Pi*SZ protein phenotypes (homozygous)] 11 μmol/L Yes No	
<ul> <li>Does the member have a diagnosis of</li> <li>A high risk AATD genetic v Yes □ No</li> <li>Other rare AATD disease-ca</li> <li>What is the member's baseline circula</li> <li>Does the member have any contraind</li> <li>Does the member have a diagnosis of</li> <li>Forced expiratory volume in</li> <li>FEV1 from &gt; 65% to &lt; 80% &gt; 100 mL/year</li> <li>Will optimal conventional treatment is</li> </ul>	ongenital deficiency of A-1 f f congenital alpha-1-antitryp variant [e.g., Pi*ZZ, Pi*Z(nu ausing alleles associated with ating serum concentration of lications to therapy? $\Box$ Yes f emphysema confirmed by a n one second (FEV1) from $\geq$ of predicted, post-bronchodi for emphysema be continued	PI $\Box$ Other: sin deficiency (AATE ill), Pi*(null)(null), or h serum AAT level < F AATD? $\Box$ No any of the following? 30% to $\leq 65\%$ of pre- ilator, and a rapid decl d (e.g. bronchodilators	) confirmed by ONE of the following? Pi*SZ protein phenotypes (homozygous)] 11 μmol/L Yes No	
<ul> <li>Does the member have a diagnosis of</li> <li>A high risk AATD genetic v Yes □ No</li> <li>Other rare AATD disease-ca</li> <li>What is the member's baseline circula</li> <li>Does the member have any contraind</li> <li>Does the member have a diagnosis of</li> <li>Forced expiratory volume in</li> <li>FEV1 from &gt; 65% to &lt; 80% &gt; 100 mL/year</li> </ul>	ongenital deficiency of A-1 $\therefore$ f congenital alpha-1-antitryp variant [e.g., Pi*ZZ, Pi*Z(nu ausing alleles associated with ating serum concentration of lications to therapy? f emphysema confirmed by a n one second (FEV1) from $\ge$ of predicted, post-bronchodi for emphysema be continued $\therefore$ or ex-smoker? Yes	PI $\Box$ Other: sin deficiency (AATE ill), Pi*(null)(null), or h serum AAT level < F AATD? $\Box$ No any of the following? 30% to $\leq 65\%$ of pre- ilator, and a rapid decl d (e.g. bronchodilators No	<ul> <li>b) confirmed by ONE of the following?</li> <li>Pi*SZ protein phenotypes (homozygous)]</li> <li>11 μmol/L Yes No</li> <li>Hicted, post-bronchodilator</li> <li>ine in lung function showing a change in FEV1</li> <li>, supplemental oxygen, etc.)?</li> <li>Yes No</li> </ul>	
<ul> <li>Does the member have a diagnosis of</li> <li>A high risk AATD genetic v Yes □ No</li> <li>Other rare AATD disease-ca</li> <li>What is the member's baseline circula</li> <li>Does the member have any contraind</li> <li>Does the member have a diagnosis of</li> <li>Forced expiratory volume in</li> <li>FEV1 from &gt; 65% to &lt; 80% &gt; 100 mL/year</li> <li>Will optimal conventional treatment is</li> </ul>	ongenital deficiency of A-1 f congenital alpha-1-antitryp variant [e.g., Pi*ZZ, Pi*Z(nu ausing alleles associated with ating serum concentration of fications to therapy? $\Box$ Yes f emphysema confirmed by a n one second (FEV1) from $\geq$ of predicted, post-bronchodi for emphysema be continued to rex-smoker? $\Box$ Yes $\Box$ <b>CURRENT or PR</b>	PI $\Box$ Other: sin deficiency (AATI ill), Pi*(null)(null), or h serum AAT level < f AATD? $\Box$ No any of the following? 30% to $\leq 65\%$ of pre- ilator, and a rapid decl d (e.g. bronchodilators No EVIOUS THERAPY	D) confirmed by ONE of the following? Pi*SZ protein phenotypes (homozygous)] I1 μmol/L Yes No Hicted, post-bronchodilator ine in lung function showing a change in FEV1 , supplemental oxygen, etc.)? Yes No	
<ul> <li>Does the member have a diagnosis of</li> <li>A high risk AATD genetic v Yes □ No</li> <li>Other rare AATD disease-ca</li> <li>What is the member's baseline circula</li> <li>Does the member have any contraind</li> <li>Does the member have a diagnosis of</li> <li>Forced expiratory volume in</li> <li>FEV1 from &gt; 65% to &lt; 80% &gt; 100 mL/year</li> <li>Will optimal conventional treatment is</li> </ul>	ongenital deficiency of A-1 $\therefore$ f congenital alpha-1-antitryp variant [e.g., Pi*ZZ, Pi*Z(nu ausing alleles associated with ating serum concentration of lications to therapy? f emphysema confirmed by a n one second (FEV1) from $\ge$ of predicted, post-bronchodi for emphysema be continued $\therefore$ or ex-smoker? Yes	PI $\Box$ Other: sin deficiency (AATE ill), Pi*(null)(null), or h serum AAT level < F AATD? $\Box$ No any of the following? 30% to $\leq 65\%$ of pre- ilator, and a rapid decl d (e.g. bronchodilators No	<ul> <li>b) confirmed by ONE of the following?</li> <li>Pi*SZ protein phenotypes (homozygous)]</li> <li>11 μmol/L Yes No</li> <li>Hicted, post-bronchodilator</li> <li>ine in lung function showing a change in FEV1</li> <li>, supplemental oxygen, etc.)?</li> <li>Yes No</li> </ul>	
<ul> <li>Does the member have a diagnosis of</li> <li>A high risk AATD genetic v Yes □ No</li> <li>Other rare AATD disease-ca</li> <li>What is the member's baseline circula</li> <li>Does the member have any contraind</li> <li>Does the member have a diagnosis of</li> <li>Forced expiratory volume in</li> <li>FEV1 from &gt; 65% to &lt; 80% &gt; 100 mL/year</li> <li>Will optimal conventional treatment is</li> </ul>	ongenital deficiency of A-1 f congenital alpha-1-antitryp variant [e.g., Pi*ZZ, Pi*Z(nu ausing alleles associated with ating serum concentration of fications to therapy? $\Box$ Yes f emphysema confirmed by a n one second (FEV1) from $\geq$ of predicted, post-bronchodi for emphysema be continued to rex-smoker? $\Box$ Yes $\Box$ <b>CURRENT or PR</b>	PI $\Box$ Other: sin deficiency (AATI ill), Pi*(null)(null), or h serum AAT level < f AATD? $\Box$ No any of the following? 30% to $\leq 65\%$ of pre- ilator, and a rapid decl d (e.g. bronchodilators No EVIOUS THERAPY	D) confirmed by ONE of the following? Pi*SZ protein phenotypes (homozygous)] I1 μmol/L Yes No Hicted, post-bronchodilator ine in lung function showing a change in FEV1 , supplemental oxygen, etc.)? Yes No	
<ul> <li>Does the member have a diagnosis of</li> <li>A high risk AATD genetic v Yes □ No</li> <li>Other rare AATD disease-ca</li> <li>What is the member's baseline circula</li> <li>Does the member have any contraind</li> <li>Does the member have a diagnosis of</li> <li>Forced expiratory volume in</li> <li>FEV1 from &gt; 65% to &lt; 80% &gt; 100 mL/year</li> <li>Will optimal conventional treatment is</li> </ul>	ongenital deficiency of A-1 f congenital alpha-1-antitryp variant [e.g., Pi*ZZ, Pi*Z(nu ausing alleles associated with ating serum concentration of lications to therapy? ☐ Yes f emphysema confirmed by a n one second (FEV1) from ≥ of predicted, post-bronchodi for emphysema be continued to rex-smoker? ☐ Yes ☐ CURRENT or PR Strength/ Frequency	PI ☐ Other: sin deficiency (AATI II), Pi*(null)(null), or h serum AAT level < f AATD? ☐ No any of the following? 30% to ≤ 65% of pre- ilator, and a rapid decl d (e.g. bronchodilators No EVIOUS THERAPY Dates of Therapy	D) confirmed by ONE of the following? Pi*SZ protein phenotypes (homozygous)] I1 μmol/L Yes No Hicted, post-bronchodilator ine in lung function showing a change in FEV1 , supplemental oxygen, etc.)? Yes No	
Does the member have a diagnosis of A high risk AATD genetic vyyes □ No Other rare AATD disease-car What is the member's baseline circular Does the member have any contrained Does the member have any contrained Does the member have a diagnosis of Forced expiratory volume in FEV1 from > 65% to < 80% > 100 mL/year Will optimal conventional treatment Is the member currently a nonsmoker Medication Name	ongenital deficiency of A-1 f congenital alpha-1-antitryp variant [e.g., Pi*ZZ, Pi*Z(nu ausing alleles associated with ating serum concentration of lications to therapy? ☐ Yes f emphysema confirmed by a n one second (FEV1) from ≥ of predicted, post-bronchodi for emphysema be continued to rex-smoker? ☐ Yes ☐ CURRENT or PR Strength/ Frequency REAUTH	PI $\Box$ Other: sin deficiency (AATE isin deficiency (AATE ill), Pi*(null)(null), or h serum AAT level < f AATD? $\Box$ No any of the following? 30% to $\leq 65\%$ of pre- ilator, and a rapid decl d (e.g. bronchodilators No EVIOUS THERAPY Dates of Therapy ORIZATION	D) confirmed by ONE of the following? Pi*SZ protein phenotypes (homozygous)] □ 11 µmol/L □ Yes □ No dicted, post-bronchodilator ine in lung function showing a change in FEV1 , supplemental oxygen, etc.)? □ Yes □ No Status (Discontinued & Why/Current)	
Does the member have a diagnosis of A high risk AATD genetic vyyes No Other rare AATD disease-car What is the member's baseline circular Does the member have any contrained Does the member have any contrained Does the member have a diagnosis of Forced expiratory volume in FEV1 from > 65% to < 80% > 100 mL/year Will optimal conventional treatment Is the member currently a nonsmoker Medication Name	ongenital deficiency of A-1 f congenital alpha-1-antitryp variant [e.g., Pi*ZZ, Pi*Z(nu ausing alleles associated with ating serum concentration of lications to therapy? ☐ Yes f emphysema confirmed by a n one second (FEV1) from ≥ of predicted, post-bronchodi for emphysema be continued or ex-smoker? ☐ Yes ☐ CURRENT or PR Strength/ Frequency REAUTH( of the signs and symptoms i	PI $\Box$ Other: sin deficiency (AATE isin deficiency (AATE ill), Pi*(null)(null), or h serum AAT level < f AATD? $\Box$ No any of the following? 30% to $\leq 65\%$ of pre- ilator, and a rapid decl d (e.g. bronchodilators No EVIOUS THERAPY Dates of Therapy ORIZATION including slowed prog	D) confirmed by ONE of the following? Pi*SZ protein phenotypes (homozygous)] □ 11 µmol/L □ Yes □ No dicted, post-bronchodilator ine in lung function showing a change in FEV1 , supplemental oxygen, etc.)? □ Yes □ No Status (Discontinued & Why/Current) ression of emphysema as evidenced by	
Does the member have a diagnosis of A high risk AATD genetic vyyes No Other rare AATD disease-car What is the member's baseline circular Does the member have any contrained Does the member have any contrained Does the member have a diagnosis of Forced expiratory volume in FEV1 from > 65% to < 80% > 100 mL/year Will optimal conventional treatment Is the member currently a nonsmoker Medication Name Is there improvement or stabilization annual spirometry testing or a decrease	ongenital deficiency of A-1 f congenital alpha-1-antitryp variant [e.g., Pi*ZZ, Pi*Z(nu ausing alleles associated with ating serum concentration of lications to therapy? ☐ Yes f emphysema confirmed by a n one second (FEV1) from ≥ of predicted, post-bronchodi for emphysema be continued or ex-smoker? ☐ Yes ☐ CURRENT or PR Strength/ Frequency REAUTH( of the signs and symptoms i	PI $\Box$ Other: sin deficiency (AATE isin deficiency (AATE ill), Pi*(null)(null), or h serum AAT level < f AATD? $\Box$ No any of the following? 30% to $\leq 65\%$ of pre- ilator, and a rapid decl d (e.g. bronchodilators No EVIOUS THERAPY Dates of Therapy ORIZATION including slowed prog	D) confirmed by ONE of the following? Pi*SZ protein phenotypes (homozygous)] □ 11 µmol/L □ Yes □ No dicted, post-bronchodilator ine in lung function showing a change in FEV1 , supplemental oxygen, etc.)? □ Yes □ No Status (Discontinued & Why/Current) ression of emphysema as evidenced by	
Does the member have a diagnosis of A high risk AATD genetic vyyes No Other rare AATD disease-car What is the member's baseline circular Does the member have any contraind Does the member have any contraind Does the member have a diagnosis of Forced expiratory volume in FEV1 from > 65% to < 80% > 100 mL/year Will optimal conventional treatment is Is the member currently a nonsmoker Medication Name Is there improvement or stabilization annual spirometry testing or a decrease Please describe:	ongenital deficiency of A-1 f congenital alpha-1-antitryp variant [e.g., Pi*ZZ, Pi*Z(nu ausing alleles associated with ating serum concentration of fications to therapy? ☐ Yes f emphysema confirmed by a n one second (FEV1) from ≥ of predicted, post-bronchodi for emphysema be continued or ex-smoker? ☐ Yes ☐ CURRENT or PR Strength/ Frequency REAUTH( of the signs and symptoms i se in frequency, duration or a	PI $\Box$ Other: sin deficiency (AATE isin deficiency (AATE ill), Pi*(null)(null), or h serum AAT level < f AATD? $\Box$ No any of the following? 30% to $\leq 65\%$ of pre- ilator, and a rapid decl d (e.g. bronchodilators No EVIOUS THERAPY Dates of Therapy ORIZATION including slowed prog severity of exacerbati	D) confirmed by ONE of the following? Pi*SZ protein phenotypes (homozygous)] □ 11 µmol/L □ Yes □ No dicted, post-bronchodilator ine in lung function showing a change in FEV1 , supplemental oxygen, etc.)? □ Yes □ No Status (Discontinued & Why/Current) ression of emphysema as evidenced by ons? □ Yes □ No	
Does the member have a diagnosis of A high risk AATD genetic vyyes No Other rare AATD disease-car What is the member's baseline circular Does the member have any contraind Does the member have any contraind Does the member have a diagnosis of Forced expiratory volume in FEV1 from > 65% to < 80% > 100 mL/year Will optimal conventional treatment is Is the member currently a nonsmoker Medication Name Is there improvement or stabilization annual spirometry testing or a decrease Please describe:	ongenital deficiency of A-1 f congenital alpha-1-antitryp variant [e.g., Pi*ZZ, Pi*Z(nu ausing alleles associated with ating serum concentration of lications to therapy? ☐ Yes f emphysema confirmed by a n one second (FEV1) from ≥ of predicted, post-bronchodi for emphysema be continued or ex-smoker? ☐ Yes ☐ CURRENT or PR Strength/ Frequency REAUTH( of the signs and symptoms i	PI $\Box$ Other: sin deficiency (AATE isin deficiency (AATE ill), Pi*(null)(null), or h serum AAT level < f AATD? $\Box$ No any of the following? 30% to $\leq 65\%$ of pre- ilator, and a rapid decl d (e.g. bronchodilators No EVIOUS THERAPY Dates of Therapy ORIZATION including slowed prog severity of exacerbati	D) confirmed by ONE of the following? Pi*SZ protein phenotypes (homozygous)] □ 11 µmol/L □ Yes □ No dicted, post-bronchodilator ine in lung function showing a change in FEV1 , supplemental oxygen, etc.)? □ Yes □ No Status (Discontinued & Why/Current) ression of emphysema as evidenced by ons? □ Yes □ No	
Does the member have a diagnosis of A high risk AATD genetic vyyes No Other rare AATD disease-car What is the member's baseline circular Does the member have any contraind Does the member have any contraind Does the member have a diagnosis of Forced expiratory volume in FEV1 from > 65% to < 80% > 100 mL/year Will optimal conventional treatment is Is the member currently a nonsmoker Medication Name Is there improvement or stabilization annual spirometry testing or a decrease Please describe:	ongenital deficiency of A-1 f congenital alpha-1-antitryp variant [e.g., Pi*ZZ, Pi*Z(nu ausing alleles associated with ating serum concentration of fications to therapy? ☐ Yes f emphysema confirmed by a n one second (FEV1) from ≥ of predicted, post-bronchodi for emphysema be continued or ex-smoker? ☐ Yes ☐ CURRENT or PR Strength/ Frequency REAUTH( of the signs and symptoms i se in frequency, duration or a	PI $\Box$ Other: sin deficiency (AATE isin deficiency (AATE ill), Pi*(null)(null), or h serum AAT level < f AATD? $\Box$ No any of the following? 30% to $\leq 65\%$ of pre- ilator, and a rapid decl d (e.g. bronchodilators No EVIOUS THERAPY Dates of Therapy ORIZATION including slowed prog severity of exacerbati	D) confirmed by ONE of the following? Pi*SZ protein phenotypes (homozygous)] □ 11 µmol/L □ Yes □ No dicted, post-bronchodilator ine in lung function showing a change in FEV1 , supplemental oxygen, etc.)? □ Yes □ No Status (Discontinued & Why/Current) ression of emphysema as evidenced by ons? □ Yes □ No	
Does the member have a diagnosis of A high risk AATD genetic vyyes No Other rare AATD disease-car What is the member's baseline circular Does the member have any contraind Does the member have any contraind Does the member have a diagnosis of Forced expiratory volume in FEV1 from > 65% to < 80% > 100 mL/year Will optimal conventional treatment is Is the member currently a nonsmoker Medication Name Is there improvement or stabilization annual spirometry testing or a decrease Please describe:	ongenital deficiency of A-1 f congenital alpha-1-antitryp variant [e.g., Pi*ZZ, Pi*Z(nu ausing alleles associated with ating serum concentration of lications to therapy? ☐ Yes f emphysema confirmed by a n one second (FEV1) from ≥ of predicted, post-bronchodi for emphysema be continued of predicted, post-bronchodi for emphysema be continued c or ex-smoker? ☐ Yes ☐ CURRENT or PR Strength/ Frequency REAUTHO of the signs and symptoms i se in frequency, duration or a PORTING INFORMATIO	PI $\Box$ Other: sin deficiency (AATE isin deficiency (AATE ill), Pi*(null)(null), or h serum AAT level < f AATD? $\Box$ No any of the following? 30% to $\leq 65\%$ of pre- ilator, and a rapid decl d (e.g. bronchodilators No EVIOUS THERAPY Dates of Therapy ORIZATION including slowed prog severity of exacerbati	D) confirmed by ONE of the following? Pi*SZ protein phenotypes (homozygous)] □ 11 µmol/L □ Yes □ No dicted, post-bronchodilator ine in lung function showing a change in FEV1 , supplemental oxygen, etc.)? □ Yes □ No Status (Discontinued & Why/Current) ression of emphysema as evidenced by ons? □ Yes □ No	