

References

E-1

1. Agency for Healthcare Research and Quality (AHRQ).
2. American College of Occupational and Environmental Medicine (ACOEM).
3. CMS Transmittal AB-02-136, CR 2371.
4. Harvey LA, Brosseau L, Herbert RD. Continuous passive motion following total knee arthroplasty in people with arthritis. Cochrane Database of Systematic Reviews. 2014 ;(2):CD004260
5. Mittenenthal S. 5 physical therapy treatments you probably don't need: Here's what you should try instead. Consumer Reports. 2014.
6. NCD 280.1 for Durable Medical Equipment Reference List.
7. Steadman J, Briggs K, Matheny L, et al. Outcomes following microfracture of full-thickness articular cartilage lesions of the knee in adolescent patients. *J Knee Sur.* 2015; 28(02): 145-150.
8. Ruggiero L, Logerstedt D, Park M, et al. Partial weight bearing and continuous passive motion for rehabilitation following microfracture surgery: A multiscale finite element simulation. *Osteoarthr Cartilage.* 2014; 22: S107-S108.
9. Hill, A, Palmer J, Tanner S, et al. Use of continuous passive motion in the postoperative treatment of intra-articular knee fractures. *J Bone Joint Surg.* 2014; 96(14): e118.
10. Bertra Á, Duska Z, Tóth F, et al. Clinical experiences with cartilage repair techniques: Outcomes, indications, contraindications and rehabilitation. *Joint Disease and Related Surgery.* 2015; 26(2): 84-96.
11. American Academy of Orthopaedic Surgeons. Surgical management of osteoarthritis of the knee. Evidence-based clinical practice guideline. 2015.
12. American Physical Therapy Association (APTA). Five things physical therapists and patients should question. American Board of Internal Medicine (ABIM). 2014.
13. He, ML. Continuous passive motion for preventing venous thromboembolism after total knee arthroplasty (Review). Cochrane Database of Systematic Reviews. 2014; 7: 1-3.
14. Lopez-Liria, R. Home-based versus hospital-based rehabilitation program after total knee replacement. *Biomed Res Int.* 2015: 1-9.
15. Brennan, G. Outpatient rehabilitation care process factors and clinical outcomes among patients discharged home following unilateral total knee arthroplasty. *J Arthro.* 2015; 30: 885- 890.
16. The American Physical Therapy Association. Don't use continuous passive motion machines for the postoperative management of patients following uncomplicated total knee replacement. 2014.
17. Curtis Derek, Benche Jesper, Mygind Bente. The effect of training in an interactive dynamic stander on ankle dorsiflexion and gross motor function in children with cerebral palsy. *Dev Nuerorehabil.* 2014; 17(6): 393-397.