

## References

E-9

1. Mohler ER, Mondry TE. Prevention and treatment of lymphedema. UpToDate. Accessed 7/29/2015
2. Medical Advisory Committee. Position statement: Lymphedema risk reduction practices. National Lymphedema Network. May/2012. National Lymphedema Network. Accessed 7/30/2015.
3. InterQual® Level of Care Criteria 2015, Acute Care Adult, McKesson Health Solutions, LLC; 2015:INPT-9.
4. Gauglitz, GG. Management of keloid and hypertrophic scars following burn injuries. UpToDate. Last revised February 09, 2016.
5. Hochstein A. The role of compression garments in wound care and practice management. *Podiatry Management*. 2017;83-87.
6. Jin Y. Compression stockings for prevention of postthrombotic syndrome: A systematic review and meta-analysis. *Vascular and Endovascular Surgery*. 2016;50(5):328-334.
7. Kravitz S. Compression hosiery and multilayer wraps: Proper prescription and usage. *Podiatry Management*. 2017:123-130.
8. Mestre S, Calais C, Gaillard G, NouM, Pasqualini M, et al. Interest of an auto-adjustable nighttime compression sleeve (MOBIDERM Autofit) in maintenance phase of upper limb lymphedema. *Support Care Cancer*. 2017;25:2455-2462.
9. Kakkos S, Timpilis M, Patrinos P, et al. Acute effects of graduated elastic compressions stockings in patients with symptomatic varicose veins: A randomized double blind placebo controlled trial. *Eur J Vasc Endovasc Surg*. 2018;55:118-125.
10. Rabe E, Partsch H, Hafner J, et al. Indications for medical compression stockings in venous and lymphatic disorders: An evidence-based consensus statement. *Phlebology*. 2018;33(3):163-184.