

References

M-74

1. CMS Pub. 100-03, Medicare National Coverage Determinations Manual, Chapter 1, Section 190.11
2. Transmittal AB-02-180, CR 2323
3. Transmittal 90, CR 6138
4. Transmittal 1562, CR 6138
5. Desantis G, Hogan-Schlientz, Liska G, et al. STABLE results: Warfarin home monitoring achieves excellent INR control. *Am J Managed Care*. 2014;20(3):202-9.
6. Chen Q, Dong L, Dong , et al. Security and cost comparison of INR self-testing and conventional hospital INR testing in patients with mechanical heart valve replacement. *J Cardiothorac Surg*. 2015;10:4.
7. Downing A, Mortimer M, Hiers J. Impact of a pharmacist-driven warfarin management protocol on achieving therapeutic International Normalized Ratios. *American Journal of Health-System Pharmacy*. 2016;73(5):S69-S73 5p.
8. Hon H, Elmously A, Hoey B, et al. Inappropriate preinjury warfarin use in trauma patients: A call for a safety initiative. *Journal of Postgraduate Medicine*. 2016;62(2):73-79.
9. Benade E, Jacobson B, Louw S, Schapkaitz E. Validation of the CoaguChek XS international normalised ratio point-of-care analyser in patients at Charlotte Maxeke Johannesburg Academic Hospital, *South Africa*. *South African Medical Journal*. 2016;106(3):280-283.
10. Hue MK, Welborn T, and Nagykalai Z. In adult patients on warfarin does home self-testing of prothrombin time and/or international normalized ratio provide the same outcomes compared to testing by a home health nurse or in a clinical setting? *J Okla State Med Assoc*. 2016;109(3): 99-100.
11. Phibbs C, Love S, Jacobson A, et al. At-home versus in-clinic INR monitoring: A cost–utility analysis from the home INR study (THINRS). *J Gen Intern Med*. 2016; 31(9):1061–1067.
12. Dionizovik-Dimanovski M, Levin, A, Fried, J, et al. Correlation between home INR and core laboratory INR in patients supported with Continuous-Flow Left Ventricular Assist Devices. *ASAIO Journal*. 2015;61(4):386–390.
13. Cayley WE Jr. Self-Monitoring and Self-Management of Oral Anticoagulation. *American Family Physician*. 2017;95(11):700-701
14. Heu MK, Welborn T, Nagykalai Z. Clinical Question: In adult patients on warfarin, does-home-self-testing of prothrombin time and/or international normalized ratio provide the same outcomes compared to testing by a home health nurse or in a clinical setting? *The Journal of The Oklahoma State Medical Association*. 2016;109(3):99-100.
15. da Silva Saraiva S, Orsi FA, Santos MP, et al. Home management of INR in the public health system: feasibility of self-management of oral anticoagulation and long-term performance of individual POC devices in determining INR. *Journal of Thrombosis and Thrombolysis*. 2016;42(1):146-153.