

References

E-47

1. Malmjö, M, Huddleston E, Martin R. Biological effects of a disposable, canisterless negative pressure wound therapy system. *Eplasty*. 2014;14:e15.
2. Hurd, T, Trueman P, A Rossington. Use of portable, single-use negative pressure wound therapy in home care patients with low to moderately exuding wounds: a case series. *Ostomy Wound Management*. 2014;60(3):30-36.
3. Srivastava R, Dwivedi M, Bhagat, et al. A non-randomised, controlled clinical trial of an innovative device for negative pressure wound therapy of pressure ulcers in traumatic paraplegia patients. *International Wound Journal*. 2014.
4. Marston W, Armstrong D, Reyzelman A, et al. Multicenter randomized controlled trial comparing treatment of venous leg ulcers using mechanically versus electrically powered negative pressure wound therapy. *Adv Wound Care (New Rochelle)*. 2015;4(2):75-82.
5. Garwood C. What's new in wound treatment: a critical appraisal. *Diabetes/Metabolism Res Rev*. 2016;32:268-274.
6. U.S. Food and Drug Administration. UPDATE on Serious Complications Associated with Negative Pressure Wound Therapy Systems: FDA Safety Communication. 2011
7. Qaseem A, Humphrey LL, Forcica MA, et al. Treatment of pressure ulcers: a clinical practice guideline from the American College of Physicians. *Ann Intern Med*. 2015;162(5):370-379.
8. National Institute for Health and Care Excellence (NICE). Negative Pressure Wound Therapy for the Open Abdomen . 2013.
9. National Institute for Health and Care Excellence (NICE). Diabetic Foot Problems: Prevention and Management. 2016.
10. Hayes Directory. *Negative Pressure Wound Therapy for Chronic Wounds: Home Use*. Lansdale PA: Hayes Inc. December 2016, Update December 2018. Accessed 2/4/2019.
11. Nherera L, Trueman P, Schmoeckel M, Fatoye F. Cost-effectiveness analysis of single use negative pressure wound therapy dressings (sNPWT) compared to standard of care in reducing surgical site complications (SSC) in patients undergoing coronary artery bypass grafting surgery. *J of Cardiothorac Surg*. 2018;13:103.
12. Costa M, Achten J, Bruce J, et al. Effect of Negative Pressure Wound Therapy vs Standard Wound Management on 12-month Disability Among Adults With Severe Open Fracture of the Lower Limb. The WOLLF Randomized Clinical Trial. *JAMA*. 2018;319(22):2280-2288.