

References

E-9

1. Kakkos S, Timpilis M, Patrinos P, et al. Acute effects of graduated elastic compressions stockings in patients with symptomatic varicose veins: A randomized double-blind placebo-controlled trial. *Eur J Vasc Endovasc Surg.* 2018;55:118-125.
2. Rabe E, Partsch H, Hafner J, et al. Indications for medical compression stockings in venous and lymphatic disorders: An evidence-based consensus statement. *Phlebology.* 2018;33(3):163-184.
3. Horiuchi M, Takiguchi C, Kirihara Y, Horiuchi Y. Impact of wearing graduated compression stockings on psychological and physiological responses during prolonged sitting. *Int J Environ Res Pub Health.* 2018;15:1710.
4. Woods M. Using compression garments in the management of lymphedema. *Brit J of Nsg.* 2019;28(1).
5. Marinovich r, Li Z, Tamasi T, Quinn K, Wong S, McIntyre CW. Hemodynamic response to non - pneumatic anti-shock compression garments in patients with renal dysfunction. *BMC Nephrology* 2020;21(1):15.
6. Abe K, Tsuji T, Oka A, et al. Postural differences in the immediate effects of active exercise with compression therapy on lower limb lymphedema. *Support Care Cancer.* 2021;29(11):6535-6543.
7. Rabe E, Partsch H, Morrison N, et al. Risks and contraindications of medical compression treatment - A critical reappraisal. An international consensus statement. *Phlebology.* 2020;35(7):447-460.
8. Moiemmen N, Mathers J, Jones L, et al. Pressure garment to prevent abnormal scarring after burn injury in adults and children: the PEGASUS feasibility RCT and mixed-methods study. *Health Technol Assess.* 2018;22(36):1-162.
9. Wiseman J, Ware RS, Simons M, et al. Effectiveness of topical silicone gel and pressure garment therapy for burn scar prevention and management in children: a randomized controlled trial. *Clin Rehabil.* 2020;34(1):120-131.
10. Xiong Y, Tao X. Compression garments for medical therapy and sports. *Polymers (Basel).* 2018;10(6):663.
11. DeBruler DM, Baumann ME, Zbinden JC, et al. Improved scar outcomes with increased daily duration of pressure garment therapy. *Adv Wound Care (New Rochelle).* 2020;9(8):453-461.
12. Tantawy SA, Abdelbasset WK, Nambi G, Kamel DM. Comparative study between the effects of kinesiio taping and pressure garment on secondary upper extremity lymphedema and quality of life following mastectomy: A randomized controlled trial. *Integr Cancer Ther.* 2019;18:1534735419847276.
13. Chitambira, F. Patient perspectives: Explaining low rates of compliance to compression therapy. *Wound Practice & Research.* 2019;27(4):168-174.