

## **References**

**L-30**

1. Chan R. Biochemical markers of spontaneous preterm birth in asymptomatic women. *BioMed Research International*. 2014;2014:164081.
2. Farag A, Mohammed M, Ellaithy M, Salama H. et al. Blind vaginal fetal fibronectin swab for prediction of preterm birth. *Obstet Gynaecol Res*. 2015;41(7):1009-1017.
3. Hadži-Lega M, Markova A, Stefanovic M, and Tantrovski M. Interleukin 6 and fetal fibronectin as predictors of preterm delivery in symptomatic patients. *Bosn Basic MedSci*. 2015;15(1):51-56.
4. van Barren G-J, et al. Predictive value of cervical length measurement and fibronectin testing in threatened preterm labor. *Obstetrics & Gynecology*. 2014;123(6):1185-1193.
5. Esplin M, Elovitz M, Reddy U, et al. Predictive accuracy of serial transvaginal cervical lengths and quantitative vaginal fetal fibronectin levels for spontaneous preterm birth among nulliparous women. *JAMA*. 2017;317(10):1047-1056.
6. Boots AB, Sanchez-Ramos L, Bowers DM, Kaunitz AM, Zamora J, Schlattmann P. The shortterm prediction of preterm birth: a systematic review and diagnostic metaanalysis. *Am J Obstet and Gynecol*. 2014;210(1):54-e1.
7. NICE guidelines [NG25: 1.7.5]. Preterm labour and birth. Publish date: November 2015.
8. The Society for Maternal Fetal Medicine. When to use fetal fibronectin. Last reaffirmed: August 1, 2016.
9. Hayes, Inc. Hayes Clinical Research Response. *Tests for Premature Rupture of Fetal Membranes*. Lansdale, PA: Hayes, Inc.; December 26, 2017. Accessed January 3, 2019.
10. Ruma M, Bittner K, Soh C. Current perspectives on the use of fetal fibronectin testing in preterm labor diagnosis and management. *AM J Manag Care*. 2017;23:-S0. Accessed January 4, 2019.