

References

M-34

1. InterQual® Level of Care Criteria. Acute Care Adult. 2017.
2. H. Singh, R. J. Cooper, C.Wai Lee, et al. Mapping cortical haemodynamics during neonatal seizures using diffuse optical tomography: a case study. *NeuroImage: Clinical*. 2014;256-265.
3. V. S. Wasade , K. Elisevich, R. Tahir, et.al. Long-term seizure and psychosocial outcomes after resective surgery for intractable epilepsy. *Epilepsy & behavior*. 2015;43:122-127.
4. Coito, A. Altered directed functional connectivity in temporal lobe epilepsy in the absence of interictal spikes: A high density EEG study. *Epilepsia*. 2016;57(3):402–411.
5. Vargas, R. Sleep electroencephalography and heart rate variability interdependence amongst healthy subjects and insomnia/schizophrenia patients. *Med Biol Eng Comput*. 2016;54:77–91.
6. Acharya, J. American Clinical Neurophysiology Society Guideline 3: A Proposal for Standard Montages to Be Used in Clinical EEG. *J Clin Neurophysiol*. 2016;33(4):312-316.
9. Hayes, Inc. Hayes Medical Technology Directory Report. *Video Electronencephalogram (VEEG) for Diagnosis and Management of Epilepsy in Adults*. Lansdale, PA: Hayes, Inc.; October, 2017. Accessed August 24, 2018.
8. Fahimi G, Tabatabaei S, Fahimi E, Rajebi H. Index of Theta/Alpha Ratio of the Quantitative Electroencephalogram in Alzheimer's Disease: A Case-Control Study. *Acta Medica Iranica*. August 2017;55(8):502-506. Accessed August 27, 2018.