

References

O-24

1. Sutdet P, Polhan N, Prakotmongkol V, et al. The effect of tibia inclination on gait pattern of stroke patients with genu recurvatum who wear an Ankle Foot Orthosis: A pilot study. *Tech & Disabil.* 2019;31(3):77-82.
2. Barkocy M, Schilz J, Heimerl S, et al. The effectiveness of serial casting and ankle foot orthoses in treating toe walking in children with autism spectrum disorder. *Pediatr Phys Ther.* 2021;33(2):83-90.
3. Choo YJ, Chang MC. Effectiveness of an ankle-foot orthosis on walking in patients with stroke: A systematic review and meta-analysis. *Sci Rep.* 2021;11(1):15879.
4. Daryabor A, Yamamoto S, Orendurff M, Kobayashi T. Effect of types of ankle-foot orthoses on energy expenditure metrics during walking in individuals with stroke: A systematic review. *Disabil Rehabil.* 2022;44(2):166-176.
5. Gerez LF, Vieira AFC. Design of an adjustable stance-control knee-ankle-foot orthosis for pediatric population. *J Pediatr Rehabil Med.* 2019;12(3):305-312.
6. Johnston TE, Keller S, Denzer-Weiler C, Brown L. A Clinical practice guideline for the use of ankle-foot orthoses and functional electrical stimulation post-stroke. *J Neurol Phys Ther.* 2021;45(2):112-196.
7. Ounpuu S, Garibay E, Acsadi G, et al. The impact of orthoses on gait in children with Charcot-Marie-Tooth disease. *Gait Posture.* 2021;85:198-204.
8. Schwarze M, Bartsch LP, Block J, et al. A comparison between laterally wedged insoles and ankle-foot orthoses for the treatment of medial osteoarthritis of the knee: A randomized cross-over trial. *Clin Rehabil.* 2021;35(7):1032-1043.