

References

P-1

1. Standard of Care for Microvascular Complications and Foot Care. American Diabetes Association (ADA). *Diabetes Care*. 2015;38(Suppl1):S58-S66.
2. McCulloch DK. Overview of medical care in adults with diabetes mellitus. UpToDate. Last updated December 16, 2015.
3. Wilson O, Kirwan J, Dures, E. et al. The experience of foot problems and decisions to access foot care in patients with rheumatoid arthritis: A qualitative study. *J Foot Ankle Res*. 2017;10:4.
4. Allen M, Does A, Gunst C. Improving diabetic foot screening at a primary care clinic: A quality improvement project. *Afr J Prim Health Care Fam Med*. 2016;8(1):955.
5. Mills J. Clinical practice recommendations for management of the diabetic foot in the wound clinic. *Today's Wound Clinic*. 2016;10(6):16-18.
6. DeBrule, M. Operative treatment of ingrown toenail by nail fold resection without matricectomy. *J Am Podiatr Med Assoc*. 2015;105(4):295-301.
7. Gular O, Tuna H, Erdil M, et al. Nail braces as an alternative treatment for ingrown toenails: results from a comparison with the Winograd technique. *The Journal of Foot and Ankle Surgery*. 2015;54(4):620-624.
8. Bernardshaw S.V., Sagedal L, Michelet K, Brudvik C. Postoperative treatment after partial nail ablation of ingrown toenails—does it matter what we recommend? A blinded randomized study. *Scandinavian Journal of Primary Health Care*. 2019;37(2):165–173.
9. Shemer A, Gupta A.K., Amichai B, Farhi R, Baran R, et al. An open comparative study of nail drilling as adjunctive treatment for toenail onychomycosis. *J Dermatolog Treat*. 2016;27(5):480-483.
10. Bernardshaw S.V., Safedal L, Michelet K, Brudvik. Postoperative treatment after partial nail ablation of ingrown toenails —does it matter what we recommend? A blinded randomised study. *Scand J Prim Health Care*. 2019;37(2):165-173.
11. Dars S, Banwell H.A., Mariccianni L. The use of urea for the treatment of onychomycosis: a systematic review. *J Foot Ankle Res*. 2019;12(22):1-11.