

## References

S-131

1. Altomare DF, Giuratrabocchetta S, Knowles CH, et al. Long-term outcomes of sacral nerve stimulation for faecal incontinence. *Br J Surg*. 2015;102(4):407-415.
2. Duelund-Jakobsen J, Worsoe J, Lundby L, et al. Management of patients with faecal incontinence. *Therap Adv Gastroenterol*. 2016;9(1):86–97.
3. Bielefeldt K. Adverse events of sacral neuromodulation for fecal incontinence reported to the federal drug administration. *World J Gastrointest Pharmacol Ther*. 2016;7(2):294–305.
4. Amundsen CL, Richter HE, Menefee SA, et al. OnabotulinumtoxinA vs sacral neuromodulation on refractory urgency urinary incontinence in women: a randomized clinical trial. *JAMA*. 2016; 316(13):1366-1374.
5. Thaha MA, Abukar AA, Thin NN, et al. Sacral nerve stimulation for faecal incontinence and constipation in adults. *Cochrane Database Syst Rev*. 2015(8):CD004464.
6. Siegel S, Noblett K, Mangel J, et al. Results of a prospective, randomized, multicenter study evaluating sacral neuromodulation with InterStim therapy compared to standard medical therapy at 6-months in subjects with mild symptoms of overactive bladder. *Neurourol Urodyn*. 2015;34(3):224-30.
7. Gormley EA, Lightner DJ, Faraday M, et al. Diagnosis and Treatment of Overactive Bladder (Non-Neurogenic) in Adults: AUA/SUFU Guideline Amendment. *J Urol*. 2015;193(5):1572-1580.
8. National Institute for Health and Care Excellence (NICE). Faecal incontinence in adults: management [CG49]. 2007.
9. Noblett K, Siegel S, Mangel J, et al. Results of a prospective, multicenter study evaluating quality of life, safety, and efficacy of sacral neuromodulation at twelve months in subjects with symptoms of overactive bladder. *Neurourol Urodyn*. 2016; 35(2): 246-51.
10. Dinning PG, Hunt L, Patton V, et al. Treatment efficacy of sacral nerve stimulation in slow transit constipation: a two-phase, double blind randomized controlled crossover study. *Am J Gastroenterol*. 2015; 110(5):733-40.
11. Gormley EA, Lightner DJ, Faraday M, et al. Diagnosis and treatment of overactive bladder (non-neurogenic) in adults: AUA/SUFU guideline amendment. *J Urol*. 2015; 193(5):1572-80.
12. Pilkington SA, Emmett C, Knowles CH, et al. Surgery for constipation: Systematic review and practice recommendations: Results V: Sacral nerve stimulation. *Colorectal Dis*. 2017; 19(3):92-100.
13. Hayes, Inc. Hayes Health Technology Assessment. *Staged approach to sacral nerve stimulation for treatment of fecal incontinence*. Lansdale, PA: Hayes, Inc.; Published on 12/29/2018. Reviewed on 12/13/2018. Accessed on 12/26/2019.