

References

S-137

1. Dobrinja C, Bernardi S, Fabris B, et al. Surgical and Pathological Changes after Radiofrequency Ablation of Thyroid Nodules. *International Journal of Endocrinology*. 2015;2015:576576.
2. Howington JA, Blum MG, Chang AC, et al. Treatment of stage I and II non-small cell lung cancer: Diagnosis and management of lung cancer, 3rd ed: American College of Chest Physicians evidence-based clinical practice guidelines. *Chest*. 2013;143(5 Suppl):e278S-313S.
3. National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology: Nonsmall cell lung cancer. v.7.2015.
4. National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology: Colon Cancer. v.3.2015.
5. National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology: Thyroid Carcinoma. v2.2015.
6. National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology: Kidney Cancer. v2.2017.
7. Nguyen T, Hattery E, Khatri V. Radiofrequency ablation and breast cancer: A review. *Gland Surgery*. 2014 May; 3(2): 128–135.
8. Katsanos K, Mailli L, Krokidis M, et al. Systematic review and meta-analysis of thermal ablation versus surgical nephrectomy for small renal tumours. *Cardiovascular Interventional Radiology*. April 2014;37(2):427-437.
9. Shangqian W, Chao Q, Zhihang P, et al. Radiofrequency ablation versus partial nephrectomy for the treatment of clinical stage 1 renal masses: A systematic review and meta-analysis. *Chinese Medical Journal (Engl)*. Jul 2014;127(13):2497-2503.
10. Weber M, Sprengel S, Omlor G, et al. Clinical long-term outcome, technical success, and cost analysis of radiofrequency ablation for the treatment of osteoblastomas and spinal osteoid osteomas in comparison to open surgical resection. *Skeletal Radiol*. 2015; 44:981-993.
11. Chen J, Zhang C, Li F, et al. A meta-analysis of clinical trials assessing the effect of radiofrequency ablation for breast cancer. *Onco Targets Ther*. 2016;9:1759-66.
12. Valcavi R, Tsamatropoulos P. Health-related quality of life after percutaneous radiofrequency ablation of cold, solid, benign thyroid nodules; a 2 year follow-up study in 40 patients. *Endocrine practice*. 2015;021:887-896.
13. Ratko TA, Vats V, Brock J, et al. Local Nonsurgical Therapies for Stage I and Symptomatic Obstructive NonSmall-Cell Lung Cancer (Comparative Effectiveness Review No. 112). Rockville (MD): Agency for Healthcare Research and Quality; 2013.
14. National Institute for Health and Care Excellence (NICE). Percutaneous radiofrequency ablation for primary and secondary lung cancers. 2010. Accessed August 14, 2018.
15. Donington J, Ferguson M, Mazzone P, et al. American College of Chest Physicians and Society of Thoracic Surgeons consensus statement for evaluation and management for high-risk patients with stage I non-small cell lung cancer. *Chest*. Dec 2012;142(6):1620-1635.