

References

S-150

1. American College of Occupational and Environmental Medicine. Occupational Medicine Practice Guideline, 2nd Ed. 2008.
2. American Medical Association. *Current Procedural Terminology* – Professional Edition.
3. Abejon D, Garcia-del-Valle S, Fuentes M, et al. Pulsed radiofrequency in lumbar radicular pain: clinical effects in various etiological groups. *Pain Pract.* 2007 7(1):21-26.
4. Barnsley L. Percutaneous radiofrequency neurotomy for chronic neck pain: outcomes in a series of consecutive patients. *Pain Med.* 2005;6(4):282-286.
5. Boswell M, Colson J, Sehgal N, et al. A systematic review of therapeutic facet joint interventions in chronic spinal pain. *Pain Physician* 2007;10:229-253.
6. Boswell MV, Manchikanti L, Kaye AD, et al. A best-evidence systematic appraisal of the diagnostic accuracy and utility of facet (zygapophysial) joint injections in chronic spinal pain. *Pain Physician.* 2015;18:E497-E533.
7. Boswell M, Shah R, Everett C, et al. Interventional techniques in the management of chronic spinal pain: evidence-based practice guidelines. *Pain Physician.* 2005;8(1):1-47.
8. Buijs E, van Wijk R, Geurts J, Weeseman et al. Radiofrequency lumbar facet denervation: A comparative study of the reproducibility of lesion size after 2 current radiofrequency techniques. *Reg Anesth Pain Med.* 2004;29(5):400-407.
9. Cahana A, Van Zundert J, Macrea L, et al. Pulsed radiofrequency: current clinical and biological literature available. *Pain Med.* 2006 t;7(5):411-423.
10. Civelek E, Cansever T, Kabatas S, et al. Comparison of effectiveness of facet joint injection and radiofrequency denervation in chronic low back pain. *Turk Neurosurg.* 2012; 22:200-206.
11. Cohen S, Hurley R, Buckenmaier C, et. al. Randomized placebo-controlled study evaluating lateral branch radiofrequency denervation for sacroiliac joint pain. *Anesthesiology.* 2008; 109(2): 279–288.
12. Cohen S, Hurley R, Christo P, Winkley et al. Clinical predictors of success and failure for lumbar facet radiofrequency denervation. *Clin J Pain.* 2007;23:45-52.
13. Cohen S, Raja S. Pathogenesis, diagnosis, and treatment of lumbar zygapophysial (facet) joint pain. *Anesthesiology* 2007;106:591-614.
14. Cohen SP, Williams KA, Kurihara C, et al. Multicenter, randomized, comparative costeffectiveness study comparing 0, 1, and 2 diagnostic medial branch (facet joint nerve) block treatment paradigms before lumbar facet radiofrequency denervation. *Anesthesiology.* 2010; 113:395-405.
15. Conlin A, Bhogal S, Sequeira K, Teasell R. Treatment of whiplash-associated disorders -part II: Medical and surgical interventions. *Pain Res Manag.* 2005 10(1):33-40.
16. Dobrogowski J, Wrzosek A, Wordliczek J. Radiofrequency denervation with or without addition of pentoxifylline or methylprednisolone for chronic lumbar zygapophysial joint pain. *Pharmacol Rep.* 2005;57:475-480.
17. Dreyfuss P, Halbrook B, Pauza K, et al. Efficacy and validity of radiofrequency neurotomy for chronic lumbar zygapophysial joint pain, *Spine.* 2000;25(10):1270-1277.

19. Geurts J, van Wijk R, Stolker R, Groen G. Efficacy of radiofrequency procedures for the treatment of spinal pain: a systematic review of randomized clinical trials. *Reg Anesth Pain Med*. 2001; 26(5):394-400.
20. Gofeld M, Jitendra J, Faclier G. Radiofrequency denervation of the lumbar zygapophysial joints: 10-year prospective clinical audit. *Pain Physician*. 2007;10:291-300.
21. Haldeman S, Carroll L, Cassidy JD, et al. Bone and Joint Decade 2000-2010 Task Force on Neck Pain and Its Associated Disorders. The Bone and Joint Decade 2000-2010 Task Force on Neck Pain and Its Associated Disorders: executive summary. *Spine*. 2008 33(4 Suppl):S5-A7.
22. Hancock M, Maher C, Latimer J, et al. Systematic review of tests to identify the disc, SIJ or facet joint as the source of low back pain. *Eur Spine J*. 2007;16(10):1539-1550.
23. Hooten W, Martin D, Huntoon M. Radiofrequency neurotomy for low back pain: evidence-based procedural guidelines. *Pain Med*. 2005;6(2):129-138.
24. King W, Ahmed SU, Baisden J, Patel N, Kennedy DJ, MacVicar J, Duszynski B. Diagnosis and treatment of posterior sacroiliac complex pain: a systematic review with comprehensive analysis of the published data. *Pain Med* 2015 Feb; 16(2): 257.
25. Koizuka S, Saito S, Kawauchi C, et al. Percutaneous radiofrequency lumbar facet rhizotomy guided by computed tomography fluoroscopy. *J Anesth*. 2005;19(2):167-169.
26. Joo YC, Park JY, Kim KH. Comparison of alcohol ablation with repeated thermal radiofrequency ablation in medial branch neurotomy for the treatment of recurrent thoracolumbar facet joint pain. *J Anesth*. 2013; 27:390-395.
27. Lakemeier S, Lind M, Schultz W, et al. A comparison of intraarticular lumbar facet joint steroid injections and lumbar facet joint radiofrequency denervation in the treatment of low back pain: A randomized, controlled, double-blind trial. *Anesth Analg*. 2013;117:228-235.
28. Laslett M, McDonald B, Aprill C, et al. Clinical predictors of screening lumbar zygapophysial joint blocks: development of clinical prediction rules. *Spine J*. 2006 ;6(4):370-379.
29. Laslett M, Oberg B, Aprill C, McDonald B. Zygapophysial joint blocks in chronic low back pain: a test of Revel's model as a screening test. *BMC Musculoskelet Disord*. 2004;5:43.
30. Leclaire R, Fortin L, Lambert R, et al. Radiofrequency facet joint denervation in the treatment of low back pain: a placebo-controlled clinical trial to assess efficacy, *Spine*. 2001;26(13):1411-1416.
31. Lee CH, Chung CK, Kim CH. The efficacy of conventional radiofrequency denervation in patients with chronic low back pain originating from the facet joints: a meta-analysis of randomized controlled trials. *Spine J*. 2017 Nov;17(11):1770-1780.
32. Lord S, Barnsley L, Wallis B, McDonald G, Bogduk N. Percutaneous radio-frequency neurotomy for chronic cervical zygapophysial-joint pain. *N Engl J Med*. 1996; 335:1721-1726.
33. Maas ET, Ostelo RW, Niemisto L, Jousimaa J, Hurri H, Malmivaara A, van Tulder MW. Radiofrequency denervation for chronic low back pain. *Cochrane Database Syst Rev*. 2015 Oct 23;(10):CD008572. doi: 10.1002/14651858.CD008572.pub2.
34. MacVicar J, Borowczyk JM, MacVicar AM, Loughnan BM, Bogduk N. Cervical medial branch radiofrequency neurotomy in New Zealand. *Pain Med*. 2012;13:647-654.

35. MacVicar J, Kreiner DS, Duszynski B, Kennedy DJ. Appropriate use criteria for fluoroscopically guided diagnostic and therapeutic sacroiliac interventions: results from the spine intervention society convened multispecialty collaborative. *Pain Med.* 2017;18(11):2081-2095.
36. Manchikanti L, Abdi S, Atluri S, et al. An update of comprehensive evidence-based guidelines for interventional techniques of chronic spinal pain: Part II: Guidance and recommendations. *Pain Physician.* 2013;16:S49-S283.
37. Manchikanti L, Kaye AD, Boswell MV, et al. A systematic review and best evidence synthesis of the effectiveness of therapeutic facet joint interventions in managing chronic spinal pain. *Pain Physician.* 2015;18:E535-E582.
38. Manchikanti L, Cash K, Pampati V, Fellows B. Influence of psychological variables on the diagnosis of facet joint involvement in chronic spinal pain. *Pain Physician.* 2008 ;11(2):145-160.
39. Manchikanti L, Manchukonda R, Pampati V, et al. Prevalence of facet joint pain in chronic low back pain in postsurgical patients by controlled comparative local anesthetic blocks. *Arch Phys Med Rehabil.* 2007;88(4):449-455.
40. Manchikanti L, Staats P, Singh V, et al. Evidence-based practice guidelines for interventional techniques in the management of chronic spinal pain. *Pain Phys.* 2003;6:3-81.
41. Manchukonda R, Manchikanti K, Cash K, et al. Facet joint pain in chronic spinal pain: an evaluation of prevalence and false-positive rate of diagnostic blocks. *J Spinal Disord Tech.* 2007;20(7):539-545.
42. McDonald G, Lord S, Bogduk N. Long-term follow-up of patients treated with cervical radiofrequency neurotomy for chronic neck pain. *Neurosurgery* 1999 ;45(1):61-68.
43. Mikeladze G, Espinal R, Finnegan R, et al. Pulsed radiofrequency application in treatment of chronic zygapophyseal joint pain. *Spine J.* 2003;3(5):360-362.
44. Moon JY, Lee PB, Kim YC, Choi SP, Sim WS. An alternative distal approach for the lumbar medial branch radiofrequency denervation: A prospective randomized comparative study. *Anesth Analg.* 2013;116:1133-1140.
45. Nath S, Nath C, Pettersson K. Percutaneous lumbar zygapophysial (Facet) joint neurotomy using radiofrequency current, in the management of chronic low back pain: a randomized double-blind trial. *Spine.* 2008;33(12):1291-1298.
46. Niemisto L, Kalso E, Malmivaara A, et al. Radiofrequency denervation for neck and back pain. *The Cochrane Database of Systematic Reviews.* 2006; Issue 3.
47. Oh W, Shim J. A randomized controlled trial of radiofrequency denervation of the ramus communicans nerve for chronic discogenic low back pain. *Clin J Pain* 2004;20(1):55-60.
48. Patel N, Gross A, Brown L, Gekht G. A randomized, placebo-controlled study to assess the efficacy of lateral branch neurotomy for chronic sacroiliac joint pain. *Pain Medicine.* 2012;13:383-398.
49. Sanders M, Zuurmond W. Percutaneous intra-articular lumbar facet joint denervation in the treatment of low back pain: a comparison with percutaneous extra-articular lumbar facet denervation. *Pain Clinic.* 1999;11(4):329-335.
50. Sapir DA, Gorup JM. Radiofrequency medial branch neurotomy in litigant and non-litigant patients with cervical whiplash. *Spine (Phila Pa 1976).* 2001;26:E268-E273.

51. Schofferman J, Kine G. Effectiveness of repeated radiofrequency neurotomy for lumbar facet pain. *Spine*. 2004;29(21):2471-2473.
52. Slipman C, Bhat A, Gilchrist R, et al. A critical review of the evidence for the use of zygapophysial injections and radiofrequency denervation in the treatment of low back pain, *Spine J*. 2003;3(4):310-316.
53. Speldewinde GC. Outcomes of percutaneous zygapophysial and sacroiliac joint neurotomy in a community setting. *Pain Med*. 2011;12:209-218.
54. Tekin I, Mirzai H, Ok G, Erbuyun et al. A comparison of conventional and pulsed radiofrequency denervation in the treatment of chronic facet joint pain. *Clin J Pain*. 2007;23(6):524-529.
55. Tzaan W, Tasker R. Percutaneous radiofrequency facet rhizotomy--experience with 118 procedures and reappraisal of its value. *Canadian Journal of Neurological Sciences*. 2000;27(2):125- 130.
56. Vallejo R, Benyamin R, Kramer J, et al. Pulsed radiofrequency denervation for the treatment of sacroiliac joint syndrome. *Pain Med*. 2006;7(5):429-434.
57. Van Kleef M, Barendse G, Kessels A, et al. Randomized trial of radiofrequency lumbar facet denervation for chronic low back pain. *Spine*. 1999;24(18):1937-1942.
58. Van Wijk R, Geurts J, Wynne H, et al. Radiofrequency denervation of lumbar facet joints in the treatment of chronic low back pain: a randomized, double-blind, sham lesion-controlled trial. *Clin J Pain*. 2005; 21(4):335-344.
59. Van Zundert J, Patihh J, Kessels A, et al. Pulsed radiofrequency adjacent to the cervical dorsal root ganglion in chronic cervical radicular pain: a double-blind sham controlled randomized clinical trial. *Pain*. 2007;127:173-182
60. Washington State Department of Labor and Industries. Guideline on diagnostic facet medial nerve branch blocks and facet neurotomy. *Provider Bull*. 2005;(PB 05-11):1-6.
61. Workloss Data Institute. Official Disability Guidelines 2008.