

References

S-178

1. Hyun M, Son I, Hong C, et al. Efficacy and safety of topical glycopyrrolate in patients with facial hyperhidrosis: a randomized, multicentre, double-blinded, placebo-controlled, split-face study. *J Eur Acad Dermatol Venereol*. 2015; 29(2):278-282.
2. Rieger R, Pedevilla S, Lausecker J. Quality of life after endoscopic lumbar sympathectomy for primary plantar hyperhidrosis. *World J Surg*. 2015; 39 (4): 905-911.
3. InterQual® Level of Care Criteria 2015. *Acute Care Adult*. McKesson Health Solutions, LLC.
4. Smith CC, Praiser D. Primary Focal Hyperhidrosis. *Up-to-Date*. Last reviewed April 2016.
5. Grabell DA, Hebert AA. Current and Emerging Medical Therapies for Primary Hyperhidrosis. *Dermatol Ther (Heidelb)*. 2017; 7:25-36.
6. Brehmer F, Lockmann A, Grönemeyer LL, Kretschmer L, Schön M, et al. Repetitive injections of botulinum toxin A continuously increase the duration of efficacy in primary axillary hyperhidrosis: A retrospective analysis in 101 patients. *JGSD*. 2015. Doi: 10.1111/ddg.12623.
7. Youssef T, Soliman M. Unilateral Sequential Endoscopic Thoracic Sympathectomy for Palmar Hyperhidrosis: A Proposed Technique to Overcome Compensatory Hyperhidrosis and Improve Plantar Hyperhidrosis. *J Laparoendosc Adv Surg Tech A*. 2015; 25(5): 370-374.
8. Eriksson Mirkovic S, Rystedt A, Balling M, Swartling C. Hyperhidrosis Substantially Reduces Quality of Life in Children: A Retrospective Study Describing Symptoms, Consequences and Treatment with Botulinum Toxin. *Acta Derm Venereol*. 2018; 98:103–107.
9. Wade R, Rice S, Llewellyn A, et al. Intervention for hyperhidrosis in secondary care: a systemic review and value-of-information analysis. *Health Technol Assess*. 2017; 21(80):1-280.
10. An JS, Won CH, Han JS, Park HS, Seo KK. Comparison of onabotulinumtoxinA and rimabotulinumtoxinB for the treatment of axillary hyperhidrosis. *Dermatol Surg*. 2015;41(8):960-967.
11. Hsu TH, Chen YT, Tu YK, Li CN. A systematic review of microwave-based therapy for axillary hyperhidrosis. *J Cosmet Laser Ther*. 2017;19(5):275-282.
12. Lembranca L, Wolosker N, de Campos JRM, Kauffman P, Teivelis MP, Puech-Leao P. Videothoracoscopic sympathectomy results after oxybutynin chloride treatment failure. *Ann Vasc Surg*. 2017;43:283-287.
13. de Campos JRM, Lembranca L, Fukuda JM, et al. Evaluation of patients who underwent resympathectomy for treatment of primary hyperhidrosis. *Interact Cardiovasc Thorac Surg*. 2017; 25(5):716-719.
14. Fukuda JM, Varella AYM, Teivelis MP, et al. Video-Assisted thoracoscopic sympathectomy for facial hyperhidrosis: the influence of the main site of complaint. *Ann Vasc Surg*. 2018;46:337-344.
15. Li C, Wu F, Zhang Q, Gao Q, Shi Z, Li L. Interventions for the treatment of Frey's syndrome. *Cochrane Database Syst Rev*. 2015; 3(3):CD009959.
16. Hayes Inc. Health Technology Report. *Tap Water Iontophoresis for the Treatment of Primary Hyperhidrosis*. Lansdale, Pa: Hayes, Inc.; Published: April 27, 2018. Accessed April 30, 2019.
17. National Institute of Health and Care Excellence. Endoscopic thoracic sympathectomy for primary hyperhidrosis of the upper limb. 2014. Accessed May 15, 2019.
18. National Institute of Health and Care Excellence (NICE). Endoscopic thoracic sympathectomy for primary facial blushing. 2014. Accessed May 15, 2019.

19. Sammons J, Khachemoune A. Axillary hyperhidrosis: a focused review. *J Dermatolog Treat.* 2017;28(7):582-590.

26. Doolittle J, Walker P, Mills T, Thurston J. Hyperhidrosis: an update on prevalence and severity in the United State. *Arch Dermatol Res.* 2016;308:743–749.

27. Huang L, Jiang H, Wei D, Xue Q, Ding Q, Hu R. A comparative study of thoracoscopic sympathectomy for the treatment of hand sweating. *J Thorac Dis.* 2019;11(8):3336-3340