

## References

S-181

1. InterQual® Level of Care Criteria 2019. Acute Care Adult. McKesson Health Solutions, LLC.
2. Testa L. Unprotected left main revascularization: Percutaneous coronary intervention versus coronary artery bypass. An updated systematic review and meta-analysis of randomised controlled trials. *PLOS*. 2017;1-12.
3. Patel M. ACC/AATS/AHA/ASE/ASNC/SCAI/SCCT/STS 2017 appropriate use criteria for coronary revascularization in patients with stable ischemic heart disease. *J Am Coll Card*. 2017;69(17):2212-2241.
4. Lapham RJ, Simons M. Transmyocardial laser revascularization for management of refractory angina. In: Saperia GM, ed. *UpToDate*. Waltham, MA: UpToDate Inc;2017.
5. Ramanathan K, Abel JG, Park JE et al. Surgical versus percutaneous coronary revascularization in patients with diabetes and acute coronary syndromes. *J Am Coll Cardiol*. 2017;70(24):2995-3006.
6. Esper RB, Farkouh ME, Ribeiro EE, et al.. Syntax score in patients with diabetes undergoing coronary revascularization in the Freedom trial. *J Am Coll Cardiol*. 2018;11;72(23 Part A):2826-37.
7. Lobo AS, Cantu SM, Sharkey SW, et al. Revascularization in patients with spontaneous coronary artery dissection and ST-segment elevation myocardial infarction. *JACC*. 2019;74(10):1290-300.