

## References

S-258

1. Amann W, Berg P, Gersbach P, et al; European Peripheral Vascular Disease Outcome Study SCS- EPOS. Spinal cord stimulation in the treatment of non-reconstructable stable critical leg ischaemia: results of the European Peripheral Vascular Disease Outcome Study (SCS-EPOS). *Eur J Vasc Endovasc Surg.* 2003;26(3):280-286.
2. American College of Occupational and Environmental Medicine. Occupational Medicine Practice Guideline, 2nd Ed. 2008.
3. American Medical Association. Current Procedural Terminology – 2014 Professional Edition.
4. Burns B, Watkins L, Goadsby PJ. Treatment of intractable chronic cluster headache by occipital nerve stimulation in 14 patients. *Neurology* 2009; 72: 341–345.
5. Burns B, Watkins L, Goadsby PJ. Treatment of hemicrania continua by occipital nerve stimulation with a bion device: long-term follow-up of a crossover study. *Lancet Neurol* 2008;7:1001–12.
6. Claeys L, Berg W, Jonas S. Spinal cord stimulation in the treatment of chronic critical limb ischemia. *Acta Neurochir Suppl.* 2007;97(Pt 1):259-265.
7. Cruccu G, Aziz T, Garcia-Larrea L, et al. EFNS guidelines on neurostimulation therapy for neuropathic pain. *Eur J Neurol.* 2007;14(9):952-970.
8. Daousi C, Benbow SJ, MacFarlane IA. Electrical spinal cord stimulation in the long-term treatment of chronic painful diabetic neuropathy. *Diabet Med* 2005;22:393–398.
9. de Vos CC, Meier K, Zaalberg PB, et al. Spinal cord stimulation in patients with painful diabetic neuropathy: a multicenter randomized clinical trial. *Pain* 2014;155:2426–2431.
10. Fontaine D, Christophe Sol J, Raoul S, Fabre N, Geraud G, Magne C et al. Treatment of refractory chronic cluster headache by chronic occipital nerve stimulation. *Cephalalgia* 2011;31:1101–5.
11. Forouzanfar T, Kemler M, Weber W, et al. Spinal cord stimulation in complex regional pain syndrome: cervical and lumbar devices are comparably effective. *Br J Anaesth.* 2004;92(3):348-53.
12. Gersbach P, Argitis V, Gardaz Jpet al. Late outcome of spinal cord stimulation for unreconstructable and limb-threatening lower limb ischemia. *Eur J Vasc Endovasc Surg.* 2007;33(6):717-724.
13. Harke H, Gretenkort P, Ladleif H, Rahman S. Spinal cord stimulation in sympathetically maintained complex regional pain syndrome type I with severe disability. A prospective clinical study. *Eur J Pain.* 2005;9(4):363-373.
14. Jivegård L, Augustinsson L, Holm J, et al. Effects of spinal cord stimulation (SCS) in patients with inoperable severe lower limb ischaemia: a prospective randomised controlled study. *Eur J Vasc Endovasc Surg.* 1995;9(4):421-425.
15. Kapural L, Narouze S, Janicki TI, Mekhail N. Spinal cord stimulation is an effective treatment for the chronic intractable visceral pelvic pain. *Pain Med.* 2006;7(5):440-443.
16. Kemler M, Barendse GA van Kleef M, et al. Spinal cord stimulation in patients with chronic reflex sympathetic dystrophy. *N Engl J Med.* 2000;343(9):618-624.

17. Kemler M, de Vet H, Barendse G, et al. Effect of spinal cord stimulation for chronic complex regional pain syndrome Type I: five-year final follow-up of patients in a randomized controlled trial. *J Neurosurg*. 2008;108(2):292-298.
18. Kemler M, de Vet H, Barendse G, et al. Spinal cord stimulation for chronic reflex sympathetic dystrophy – five year follow-up. *N Engl J Med*. 2006;354(22):2394-2396.
19. Kemler M, de Vet H, Barendse G, et al. The effect of spinal cord stimulation in patients with chronic reflex sympathetic dystrophy: two years' follow-up of the randomized controlled trial. *Ann Neurol*. 2004;55:13-18.
20. Kemler M, Furnee C. Economic evaluation of spinal cord stimulation for chronic reflex sympathetic dystrophy. *Neurology*. 2002;59(8):1203-1209.
21. Klomp H, Spincemaille G, Steyerberg E, et al. Spinal-cord stimulation in critical limb ischaemia: a randomised trial. ESES Study Group. *Lancet*. 1999;353(9158):1040-1044.
22. Kumar K, Hunter G, Demeria D. Spinal cord stimulation in treatment of chronic benign pain: challenges in treatment planning and present status, a 22-year experience. *Neurosurgery*. 2006;58(3):481-496; discussion 481-96.
23. Latif O, Nedeljkovic S, Stevenson L. Spinal cord stimulation for chronic intractable angina pectoris: a unified theory on its mechanism. *Clin Cardiol*. 2001;24:533-541.
24. Magis D, Allena M, Bolla M, De Pasqua V, Remacle JM, Schoenen J. Occipital nerve stimulation for drug-resistant chronic cluster headache: a prospective pilot study. *Lancet Neurol* 2007; 6: 314–321.
25. Magis D, Gerardy PY, Remacle JM, Schoenen J. Sustained effectiveness of occipital nerve stimulation in drug-resistant chronic cluster headache. *Headache* 2011;51:1191–201.
26. Magis D, Schoenen J. Advances and challenges in neurostimulation for headaches. *Lancet Neurol* 2012;11:708–19.
27. Lipton R, Goadsby P, Cady R, Aurora SK, Grosberg B, Freitag F et al. PRISM study: occipital nerve stimulation for treatment-refractory migraine. *Cephalalgia* 2009;29:30.
28. Mailis-Gagnon A, Furlan A, Sandoval J, Taylor R, Spinal cord stimulation for chronic pain.
29. Cochrane Database Syst Rev. 2004;3:CD003783.
30. Mannheimer C, Eliasson T, Augustinsson LE, et al. Electrical stimulation versus coronary artery bypass surgery in severe angina pectoris: the ESBY study. *Circulation*. 1998;97:1157-1163.
31. Marin JC, Goadsby P. Response of SUNCT (short-lasting unilateral neuralgiform headaches with conjunctival injection and tearing), SUNA (short-lasting unilateral neuralgiform headaches with autonomic symptoms) and primary stabbing headaches to occipital nerve stimulation (ONS). *Neurology* 2010;74:P04.006, abstract.
32. McGreevy K, Williams K. Contemporary insights into painful diabetic neuropathy and treatment with spinal cord stimulation. *Curr Pain Headache Rep* 2012;16:43–49.
33. National Institute for Health and Care Excellence (NICE). Pain (chronic neuropathic or ischaemic) - spinal cord stimulation: final appraisal determination. September 2008.

34. North R, Calkins S, Campbell D, et al. Automated, patient-interactive, spinal cord stimulator adjustment: a randomized controlled trial. *Neurosurgery*. 2003;52(3):572-580.
35. North R, Kidd D, Farrokhi F, Piantadosi S. Spinal cord stimulation versus repeated lumbosacral spine surgery for chronic pain: a randomized, controlled trial. *Neurosurgery*. 2005;56(1):98-106.
36. North R, Kidd D, Lee M, et al. A prospective, randomized study of spinal cord stimulation versus reoperation for failed back surgery syndrome: initial results. *Stereotact Funct Neurosurg*. 1994;62:267-272.
37. North R, Kidd H, Piantadosi S. Spinal cord stimulation versus reoperation for failed back surgery syndrome: a prospective, randomized study design. *Acta Neurochir*. 1995;64(Suppl):106-8.
38. North R, Kidd D, Shipley J, Taylor R. Spinal cord stimulation versus reoperation for failed back surgery syndrome: a cost effectiveness and cost utility analysis based on a randomized, controlled trial. *Neurosurgery*. 2007;61(2):361-368.
39. Pedrini L, Magnoni F. Spinal cord stimulation for lower limb ischemic pain treatment. Interact Page 11 of 12. *Cardiovasc Thorac Surg*. 2007;6(4):495-500.
40. Reed K. Peripheral neuromodulation and headaches: history, clinical approach, and considerations on underlying mechanisms. *Curr pain Headache Rep* 2013;17:305.
41. Reed KL, Black SB, Banta CJ 2nd, Will KR. Combined occipital and supraorbital neurostimulation for the treatment of chronic migraine headaches: initial experience. *Cephalalgia* 2010;30:260–71.
42. Saper JR, Dodick DW, Silberstein SD, McCarville S, Sun M, Goadsby PJ. Occipital nerve stimulation for the treatment of intractable chronic migraine headache: ONSTIM feasibility study. *Cephalalgia* 2011; 31:271–85.
43. Schwedt TJ, Dodick DW, Trentman TL, Zimmerman RS. Response to occipital nerve block is not useful in predicting efficacy of occipital nerve stimulation. *Cephalalgia* 2007; 27: 271–274.
44. Shanahan P, Watkins L, Matharu M. Treatment of medically intractable short--- lasting unilateral neuralgiform headache attacks with conjunctival injection and tearing (SUNCT) and short---lasting unilateral neuralgiform headache attacks with autonomic symptoms (SUNA) with occipital nerve stimulation (ONS) in 6 patients. *Cephalalgia* 2009;29:150.
45. Silberstein S, Dodick D, Saper J, Huh B, Slavin KV, Sharan A et al. Safety and efficacy of peripheral nerve stimulation of the occipital nerves for the management of chronic migraine: Results from a randomized, multicenter, double---blinded, controlled study. *Cephalalgia* 2012;32(16):1165–79.
46. Skaribas I, Calvillo O, Delikanaki---Skaribas E. Occipital peripheral nerve stimulation in the management of chronic intractable occipital neuralgia in a patient with neurofibromatosis type 1: a case report. *Journal of Medical Case Reports* 2011;5:174.Lambru G, Matharu M. Occipital nerve stimulation in primary headache syndromes. *Ther Adv Neurol Disord* 2012;5(1):57---67.
47. Slangen R, Pluijms WA, Faber CG, Dirksen CD, Kessels AG, van Kleef M. Sustained effect of spinal cord stimulation on pain and quality of life in painful diabetic peripheral neuropathy. *Br J Anaesth* 2013;111:1030– 1031.
48. Slangen R, Schaper NC, Faber CG, et al. Spinal cord stimulation and pain relief in painful diabetic peripheral neuropathy: a prospective two---center randomized controlled trial. *Diabetes Care* 2014;37:3016–3024.

49. Stancák A, Kozák J, Vrba I, et al. Functional magnetic resonance imaging of cerebral activation during spinal cord stimulation in failed back surgery syndrome patients. *Eur J Pain*. 2008;12(2):137-48.
50. Stanton-Hicks M, Burton A, Bruehl S, et al. An updated interdisciplinary clinical pathway for CRPS: report of an expert panel. *Pain Pract*. 2005; 84(3):S4-S16.
51. Sundaraj S, Johnstone C, Noore F, et al. Spinal cord stimulation: a seven-year audit. *J Clin Neurosci*.
52. 2005 Apr;12(3):264-270.
53. Taylor R, Buyten J, Buchser E. Spinal cord stimulation for complex regional pain syndrome: A systematic review of the clinical and cost-effectiveness literature and assessment of prognostic factors. *Eur J Pain*. 2006;10(2):91-101.
54. Taylor R, Taylor R. Spinal cord stimulation for failed back surgery syndrome: a decision-analytic model and cost-effectiveness analysis. *Int J Technol Assess Health Care*. 2005;21(3):351-358.
55. TenVaarwerk I, Jessurun G, DeJongste M, et al. Clinical outcome of patients treated with spinal cord stimulation for therapeutically refractory angina pectoris. The Working Group on Neurocardiology. *Heart*. 1999;82(1):82-88.
56. Tesfaye S, Watt J, Benbow S, et al. Electrical spinal-cord stimulation for painful diabetic peripheral neuropathy. *Lancet*. 1996;348:1698-1701.
57. Ubbink D, Vermeulen H. Spinal cord stimulation for critical leg ischemia: a review of effectiveness and optimal patient selection. *J Pain Symptom Manage*. 2006;31:S30-S35.  
Page 12 of 12
58. Ubbink D, Vermeulen H. Spinal cord stimulation for non-reconstructable chronic critical leg ischaemia. *Cochrane Database Syst Rev*. 2005;CD004001.
59. van Beek M, Slangen R, Schaper N, Faber C, Joosten E, Dirksen C, van Dongen R, Kessels A, van Kleef M. Sustained treatment effect of spinal cord stimulation in painful diabetic peripheral neuropathy: 24-month follow-up of a prospective two-center randomized controlled trial. *Diabetes Care* 2015;38:e132--e134.
60. Weiner RL, Reed KL (1999) Peripheral neurostimulation for control of intractable occipital neuralgia. *Neuromodulation* 2:217–22.
61. Workloss Data Institute. Official Disability Guidelines 2008.