

References

S-82

1. InterQual® Level of Care Criteria 2016. Acute Care Adult. McKesson Health Solutions, LLC.
2. Nam J, Jing H, O'Reilly D. Intra-arterial thrombolysis vs. standard treatment or intravenous thrombolysis in adults with acute ischemic stroke: a systematic review and meta-analysis. *Int Journal of Stroke*. 2013.
3. Spreen M, Martens J, Hansen B, et al. Percutaneous transluminal angioplasty and drug-eluting stents for infrapopliteal lesions in critical limb ischemia (PADI) trial. *Circ Cardiovasc Interv*. 2016; 9(2): e002376.
4. Chen I, Lee C, Chao T, et al. Impact of routine coronary catheterization in low extremity artery disease undergoing percutaneous transluminal angioplasty: Study protocol for a multi-center randomized controlled trial. *Trials*. 2016;17:112.
5. Herman M, Gornik H, Barrett C, et al. 2016 American Heart Association (AHA)/American College of Cardiology (ACC) Guideline on the management of patients with lower extremity peripheral artery disease: Executive summary. A report of the ACC/AHA task force on clinical practice guidelines. *Circulation*. 2016.
6. Wittens C, Davies A, Baekgaard N, et al. Management of chronic venous disease: Clinical practice guidelines of the European Society for Vascular Surgery (ESVS). *Eur J Vasc Endovasc Surg*. 2015;49:678-737.
7. Gray K, Beckord B, Moazzez A, et al. A Comparative Analysis of Open versus Endovascular Techniques for Management of Non-Aortic Cervicothoracic Arterial Injuries. *The American Surgeon*. 2017;83:1054-58
8. Hoving J, Marquering H, Majoie C. Endovascular treatment in patient with carotid artery dissection and intracranial occlusion: a systematic review. *Neuroradiology*. 2017;59:641-647