

## References

Y-1

1. Rahmann AE, Brauer SG, Nitz JC. A specific inpatient aquatic physiotherapy program improves strength after total hip or knee replacement surgery: A randomized controlled trial. *Arch Phys Med Rehabil.* 2009;90(5):745-755.
2. Hillier S, McIntyre A, Plummer L. Aquatic physical therapy for children with developmental coordination disorder: A pilot randomized controlled trial. *Phys Occup Ther Pediatr.* 2010;30(2):111-124.
3. Tinti G, Somera R Jr, Valente FM, Domingos CR. Benefits of kinesiotherapy and aquatic rehabilitation on sickle cell anemia. A case report. *Genet Mol Res.* 2010;9(1):360-364.
4. Lau CK, Lam KS, Siu E, et al. Physiotherapist-designed aquatic exercise programme for community-dwelling elders with osteoarthritis of the knee: A Hong Kong pilot study. *Hong Kong Med J.* 2014 Feb;20(1):16-23.
5. Weintraub M, Herrmann D, Smith G, Backonja M, Cold S. Pulsed Electromagnetic Fields to Reduce Diabetic Neuropathic Pain and Stimulate Neuronal Repair: A Randomized Controlled Trial. *Arch Phys Med Rehabil.* July 2009;90:1102-1109.
6. Gulich DR. Comparison of tissue heating between manual and hands-free ultrasound techniques. *Physiother Theory Pract.* 2010 Feb;26(2):100-6.
7. McGibbon N, Benda W, Duncan B, Silkwood-Sherer D. Immediate and Long-Term Effects of Hippotherapy on Symmetry of Adductor Muscle Activity and Functional Ability in Children with Spastic Cerebral Palsy. *Arch Phys Med Rehabil.* June 2009;90: 966-974.
8. Duffy R. Low Back Pain: An Approach to Diagnosis and Management. *Primary Care: Clinics in Office Practice.* Dec. 2010;37(4).
9. Zambito A, Bianchini D, Gatti D, Rossini M, Adami S, Viapiana O. Interferential and horizontal therapies in chronic low back pain due to multiple vertebral fractures: a randomized, double blind, clinical study. *Osteoporos Int.* 2007; 18(11): 1541-5
10. Meireles SM, Jones A, Jennings F, et al. Assessment of the effectiveness of low-level laser therapy on the hands of patients with rheumatoid arthritis: A randomized double-blind controlled trial. *Clin Rheumatol.* 2010 May;29(5):501-509
11. Jang H, Lee H. Meta-analysis of pain relief effects by laser irradiation on joint areas. *Photomed Laser Surg.* 2012;30(8):405-17.
12. Tumilty S, McDonough S, Hurley DA et al. Clinical effectiveness of low-level laser therapy as an adjunct to eccentric exercise for the treatment of Achilles' tendinopathy: A randomized controlled trial. *Arch Phys Med Rehabil.* 2012;93(5):733-9.
13. Morimoto Y, Saito A, Tokuhashi Y. Low level laser therapy for sports injuries. *Laser Therapy.* 2013(1):17-20.
14. Gross AR, Dziengo S, Boers O, et al. Low level laser therapy (LLLT) for neck pain: A systematic review and meta-regression. *Open Orthoptic Journal.* 2013 September;7:396-419.

15. Warden S, Metcalf B, Kiss Z, et. al. Low-intensity pulsed ultrasound for chronic patellar tendinopathy: A randomized, double-blind, placebo-controlled trial. *Rheumatology*. 2008; 47: 467-471.