

## References

Z-7

### Deep brain stimulation

1. Charles D, Konrad PE, Neimat JS, et al. Subthalamic nucleus deep brain stimulation in early stage Parkinson's disease. *Parkinsonism Relat Disord*. 2014;20(7):731-737.
2. Kim JH, Chang WS, Jng HH, et al. Effect of subthalamic deep brain stimulation on levodopa-induced dyskinesia in Parkinson's disease. *YMJ*. 2015;56(5):1316-1321.

### Transcutaneous electrical nerve stimulation (TENS)

1. Bjerså K and Andersson T. High frequency TENS as a complement for pain relief in postoperative transition from epidural to general analgesia after pancreatic resection. *Complement Ther Clin Pract*. 2014;20(1):5-10.
2. Palmer S, Domaille M, Cramp F, et al. Transcutaneous electrical nerve stimulation as an adjunct to education and exercise for knee osteoarthritis: A randomized controlled trial. *Arthritis Care Res*. 2014;66(3):387-394
3. Johnson M. Transcutaneous electrical nerve stimulation: Review of effectiveness. *Nurs Stand*. 2014;28(40):44-53.
4. Vitalli C, Oleg C. The efficiency of transcutaneous electrical nerve stimulation in association with gabapentin in the treatment of neuropathic pain in patients with spinal cord injury. *Rom J of Neuro*. 2014;193-196.
5. Noehren B, Dailey D, Rakel B, et al. Effect of transcutaneous electrical nerve stimulation on pain, function, and quality of life in fibromyalgia: A double-blind randomized clinical trial. *J Am Phys Ther Assoc*. 2015;95(1):129-140.

### Percutaneous electrical nerve stimulation (PENS)

1. National Institute for Health and Clinical Excellence Interventional Procedure Guidance 450. NICE website. Percutaneous electrical nerve stimulation for refractory neuropathic pain.

### Vagus nerve stimulation

1. Ben-Menachem E, Revesz D, Simon BJ, et al. Surgically implanted and non-invasive vagus nerve stimulation: A review of efficacy, safety and tolerability. *Eur J of Neuro*. 2015;22(9):1260-1268.
2. Wasade VS, Schultz L, Mohanarangan K, et al. Long-term seizure and psychosocial outcomes of vagus nerve stimulation for intractable epilepsy. *E&B*. 2015;53:31-36.
3. Gaul C, Diener HC, Silver N, et al. Non-invasive vagus nerve stimulation for PREvention and Acute treatment of chronic cluster headache (PREVA): a randomised controlled study. *Cephalalgia*. 2016;36(6):534-546.
4. Miller S, Sinclair AJ, Davies B, Matharu M. Neurostimulation in the treatment of primary headaches. *practNeurol*. 2016;16(5):362-375. doi: 10.1136/practneurol-2015-001298

5. Silberstein SD, Mechtler LL, Kudrow DB, et al. Non-invasive vagus nerve stimulation for the ACute Treatment of cluster headache: findings from the randomized, double-blind, sham-controlled ACT1 study. *Headache*. 2016;56(8):1317-1332.
6. [Kinfe TM, Pinteá B, Muhammad S, Zaremba S, Roeske S, Simon BJ, Vatter H. Cervical non-invasive vagus nerve stimulation \(nVNS\) for preventive and acute treatment of episodic and chronic migraine and migraine-associated sleep disturbance: a prospective observational cohort study. \*J Headache Pain\*. 2015;16:101.](#)
7. [Silberstein SD, Calhoun AH, Lipton RB, Grosberg BM, Cady RK, Dorlas S, et al. EVENT Study Group. Chronic migraine headache prevention with noninvasive vagus nerve stimulation: The EVENT study. \*Neurology\*. 2016;87\(5\):529-38.](#)
8. [Hayes Inc. Hayes Transforming Healthcare with Evidence. GammaCore Transcutaneous Vagus Nerve Stimulator. Sept. 2018.](#)
9. [Goadsby P, DeCoo I, Silver N, et al. Non-invasive vagus nerve stimulation for the acute treatment of episodic and chronic cluster headache: A randomized, double-blind, sham-controlled ACT2 study. \*Cephalalgia\*. 2018;38\(5\):959–969](#)
10. [Tassorelli C, Grazzi L, de Tommaso M, Pierangeli G, Martelletti P, Rainero I, et al. PRESTO Study Group. Noninvasive vagus nerve stimulation as acute therapy for migraine: The randomized PRESTO study. \*Neurology\*. 2018 Jun 15.](#)

### **Occipital nerve stimulation**

1. Dodick DW, Silberstein SD, Reed KL, et al. Safety and efficacy of peripheral nerve stimulation of the occipital nerves for the management of chronic migraine: Long-term results from a randomized, multicenter, double-blinded, controlled study. *IHS*. 2014;35(4):344-358.
2. Chen YF, Bramley G, Unwin G, et al. Occipital Nerve Stimulation for Chronic Migraine- A systematic review and meta-analysis. *PLoS ONE*. 2015;10(3):1-16.

### **HF10 Therapy (Senza)**

1. Russo M, Van Buyten J-P. 10-kHz high-frequency scs therapy: A clinical summary. *Pain Med*. 2014;16(5):934-942.
2. Kapural L, Yu C, Doust MW, et al. Novel 10-kHz high-frequency therapy (HF10 Therapy) is superior to traditional low-frequency spinal cord stimulation for the treatment of chronic back and leg pain: The SENZA-RCT Randomized Controlled Trial. *JASA*. 2015;123:851-860.
3. Perruchoud C. Paraesthesia-free spinal cord stimulation: The future, or a phase? *NeuroNews*. 2016;1-2.